

## Social determinants of health and disease in companion dogs: A cohort study from the Dog Aging Project

Brianah M. McCoy<sup>1,2,\*</sup>, Layla Brassington<sup>1,2,\*</sup>, Kelly Jin<sup>3</sup>, Greer A. Dolby<sup>4</sup>, Sandi Shrager<sup>5</sup>, Devin Collins<sup>6</sup>, Matthew Dunbar<sup>7</sup>, Dog Aging Project Consortium<sup>+</sup>, Audrey Ruple<sup>8</sup>, Noah Snyder-Mackler<sup>1,2,9,+</sup>

<sup>1</sup>School of Life Sciences, Arizona State University, Tempe, AZ, USA

<sup>2</sup>Center for Evolution and Medicine, Arizona State University, Tempe, AZ, USA

<sup>3</sup>Allen Institute for Brain Science, Seattle, WA, USA

<sup>4</sup>Department of Biology, University of Alabama at Birmingham, Birmingham AL USA

<sup>5</sup>Collaborative Health Studies Coordinating Center, Department of Biostatistics, University of Washington, Seattle, WA, USA

<sup>6</sup>Department of Sociology, University of Washington, Seattle, WA, USA

<sup>7</sup>Center for Studies in Demography & Ecology, University of Washington, Seattle, WA, USA

<sup>8</sup>Department of Population Health Sciences, Virginia-Maryland College of Veterinary Medicine, Virginia Tech, Blacksburg, VA, USA

<sup>9</sup>School for Human Evolution and Social Change, Arizona State University, Tempe, AZ, USA

\* equal contribution

<sup>+</sup>correspondence: [nsnyderm@asu.edu](mailto:nsnyderm@asu.edu)

© The Author(s) 2023. Published by Oxford University Press on behalf of the Foundation for Evolution, Medicine, and Public Health.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

## Abstract

Exposure to social environmental adversity is associated with health and survival across many social species, including humans. However, little is known about how these health and mortality effects vary across the lifespan and may be differentially impacted by various components of the environment. Here, we leveraged a relatively new and powerful model for human aging, the companion dog, to investigate which components of the social environment are associated with dog health and how these associations vary across the lifespan. We drew on comprehensive survey data collected on 21,410 dogs from the Dog Aging Project and identified five factors that together explained 33.7% of the variation in a dog's social environment. Factors capturing financial and household adversity were associated with poorer health and lower physical mobility in companion dogs, while factors that captured social support, such as living with other dogs, were associated with better health when controlling for dog age and weight. Notably, the effects of each environmental component were not equal: the effect of social support was 5x stronger than financial factors. The strength of these associations depended on the age of the dog, including a stronger relationship between owner's age and the dog's health in younger as compared to older dogs. Taken together, these findings suggest the importance of income, stability, and owner's age on owner-reported health outcomes in companion dogs and point to potential behavioral and/or environmental modifiers that can be used to promote healthy aging across species.

Accepted Manuscript

## Introduction

The social environment is one of the strongest predictors of health and mortality in many mammals, where exposure to less social adversity is associated with lower risk of disease and death [1]. In humans, higher socioeconomic status (SES) can lead to access to more resources, including healthcare, and a more predictable, less stressful environment that is associated with lower morbidity and mortality [2,3]. Additionally, social isolation—another component of the social environment—has a stronger association with mortality than does heavy smoking, heavy drinking, and lack of exercise, when controlling for SES [3,4]. These sociality-health links appear to be deeply rooted in evolutionary history: having been identified in species ranging from mice to primates [5–7].

How social adversity impacts older individuals is less clear and may vary based on timing, severity, and perceived impact of exposure to adversity [8]. Environmental stability, which is composed of positive socio-economic, built, and natural environmental factors—such as higher household income, more social connections, and access to greenspace, patterns overall health and disease risk in both early and late life [9]. In humans, urban environments with high residence turnover, high pollution levels, and low economic status are associated with reduced physical mobility, higher risk of disease, and worse health outcomes [10] that can also worsen with age [11]. However, researchers are only beginning to explore how health and mortality associations with environmental factors vary across the lifespan in humans [12,13] – knowledge that is necessary to optimize the distribution of resources and interventions to those who would benefit most. This gap in our knowledge is largely due to the long lives of humans and confounding factors that may be linked to other environmental factors (e.g., smoking).

To address this gap, we can turn to animal systems with similar biology, recent co-evolutionary history, and environments to humans, but compressed into a much shorter lifespan. Companion dogs (*Canis familiaris*) offer a unique opportunity to look across the lifespan at how and when aspects of the social and physical environment may alter aging, health, and survival due to the comparatively shortened lifespan of dogs. Dogs have been shown to be a powerful comparative model for human health and aging due to our shared biology, shared environment, and significantly shorter lifespan [14–18]. Similar to humans, the major causes of death and disease in older dogs are cancer, cardiovascular incidences, and immune-mediated diseases—making them a particularly important model for understanding

the biology of aging in mammals [18–20]. Dogs are also genetically and phenotypically diverse due to hundreds of years of selective breeding [21,22], and this genetic history and architecture have also made them a powerful model for investigating the genetic factors underlying complex traits, including many human diseases—including diseases associated with aging [23,24].

Yet, despite our shared environments and passion for our companion dogs, we still know relatively little about the health impacts of the social environmental factors on companion animals, such as dogs. This gap in our knowledge is, in large part, due to a dearth of detailed social environmental data available for large samples of dogs of all ages. We therefore have a limited understanding of how lived experiences might affect health and aging in dogs, which could: (1) inform how environmental factors impact human health and aging across the lifespan, and (2) help our companion animals live longer, healthier lives.

In humans, older individuals are often more vulnerable to adverse risk exposures due to a decreased ability to recover from exposures (‘resilience’), and a limited capacity for withstanding these exposures (‘reserve’) [25,26]. For example, older individuals who experience a stressor such as infection or surgery have more risk of adverse outcomes later in life [25]. This relationship between increased risk and adverse outcomes with age extends to companion dogs, who have been shown to have poorer outcomes for relatively common conditions at older ages as compared to younger ages even when medical and surgical treatments are provided [27,28]. Additionally, in humans, social adversity such as social isolation and undernourishment early in life can have far-reaching effects on development and, subsequently, on adult health, reproduction, and survival [29–33]. It is likely that social adversity such as reduced social companionship, less time spent outdoors, and other exposures to environmental risk factors identified in human populations can impact adult dog health and aging as well. In fact, it has been shown that early life malnutrition in dog populations can lead to long-term behavioral effects, like showing increased levels of fear and aggression, in addition to causing physical health impacts in adulthood [34]. Thus, broader research into the impacts of social adversity on health and aging in companion animals is warranted.

Here, we examined factors related to the social environment of companion dogs to determine if there were associations with health and disease outcomes, as well as a measure of frailty (mobility), using data collected by the Dog Aging Project (DAP). We focused on

these outcome measures because they all exhibit age-related changes (declines in health and mobility and increases in disease occurrences) in humans and other animals. In particular, mobility is a key phenotypic signature of frailty—the age-related increase of vulnerability to adverse outcomes. Mobility is associated with overall health [35,36] and declines with age [37,38] in both human and dog populations, suggesting that it can be used as a potential biomarker for aging and age-related disease risk. We hypothesized that exposure to higher levels of social environmental adversity in early adulthood would predict poorer age-associated health outcomes in this population of dogs, and that the strength of this effect would vary across the lifespan. Specifically, we predicted that dogs living in homes with more factors associated with signatures of adverse environments (i.e., low social integration, less stable environments, and lower SES) would be less mobile, in poorer health, and have more disease than more socially advantaged dogs within the population.

## Methods

### *Study population and recruitment*

The Dog Aging Project (DAP) is a community science project that aims to understand how genes, lifestyle, and the environment influence aging and disease outcomes [39,40]. Study participants are recruited to DAP through word-of-mouth, mainstream media, and social media. For this study we drew on the first year of survey data released on May 10, 2021, from 27,541 dogs completed by owners on or before December 31, 2020. Study data were collected and managed using REDCap (Research Electronic Data Capture) electronic data capture tools hosted through the DAP [41,42].

### *Dog Aging Project survey data*

The Health and Life Experiences Survey (HLES) includes a number of small surveys that collect information about dog demographic characteristics, physical activity, environment, dog behavior, diet, medications and preventatives, health status, and owner demographic characteristics. The DAP Environment Dataset contains geographically-defined data from secondary sources pertaining to the dog's external environment. Environmental data are based on respondents' primary and (where applicable) secondary address information, provided in the HLES owner contact form. Environmental metadata was generated from the U.S. Census Bureau (2019) and the 2015-2019 American Community Survey 5-year Public

Use Microdata Samples. Respondent addresses are geocoded and linked to existing data that characterize various aspects of the dog's external environment. 56,285 Environment data records (two per dog) are included in the 2020 Curated Data Release, reflecting all study participants who became DAP pack members on or before December 31, 2020, and released on May 10, 2021 (n=27,541). We selected only the data corresponding to the primary residence information for the year 2020. We removed participants under 2 years of age in order to only include dogs that are fully grown, following the American Animal Hospital Association life stages guidelines [43]. After applying all these criteria, 21,410 dogs remained for downstream analysis. We excluded variables that were less than 98% complete and redundant variables that captured similar metrics (e.g. swimming/aerobic activity which was captured by overall activity level). Variables that were excluded due to missingness either lacked census data or the question did not apply to the owner (i.e., potential follow-up questions associated with a specific type of home heating method).

#### *Mobility metric*

Due to the limited nature of survey questions, we attempted to generate a composite metric that could roughly capture age-related decline without someone carrying out a defined physical examination of the dog. This metric was calculated by taking the z-score for each activity related variable in the HLES (activity level, average activity intensity, on leash walk frequency, aerobic activity frequency, physical games frequency, average daily activity, and on leash walk average) then assigning each dog a score which corresponded to the average of their z-score on the activity related variables.

#### *Disease instances metric*

The disease instance outcome metric was generated by summing the number of owner-reported diseases for each dog. We excluded diseases that did not have a severe effect on overall dog health as described in [44] and detailed in SI Table 5).

#### *Stability metric*

The stability metric is a combined variable that encompasses the percentage of the population in a census tract that was in the same home 1 year ago, the percent of homes that are owned by the occupant, and the percentage of the population born in the United States. Each dog received a stability score calculated from taking the z-score of the variables and averaging them. These data were collected from the U.S. Census Bureau (2019). Some of the variables

included in this metric are human-centric measures that will also capture the stability of a dog's local environment.

### *Factor analysis*

For this study, we focused our analysis on the responses to 33 survey questions in the HLES. These 33 questions were selected a priori based on their link to at least one of the axes of the social determinants of health (e.g., financial, environmental, social connection; see table S2 and methods). We then reduced the 33 survey questions into a smaller set of latent variables using exploratory factor analysis and following well-established guidelines [45]. Briefly, we first performed parallel analysis to determine the number of factors to retain then conducted factor analysis using the Psych package (version 2.0.8) in R [46]. After determining the optimal number of factors to retain, we then performed exploratory factor analysis using the default settings for the `fa()` function.

### *Modeling of health-related variables*

We modeled each of our health metrics (owner-reported health, disease instances, and mobility score) as a function of all environmental factors, age, and weight, as well as the interaction between age and each factor (SI Figure 4). We included weight in the model to control for variance in the health-related variables associated with breed ancestry, due to large inter-breed differences in dog size [20,47]. We do not have genotypic data from the mixed-breed dogs, so could not control for breed ancestry directly. Due to their distributions and variable types, each health metric required a different linear modeling framework. We used an ordinal regression to model owner-reported health, which was scored on a Likert scale from 1-6. We modeled the count-based owner-reported disease instances using a generalized linear model with a Poisson link function. Mobility was modeled using a standard linear model. For all outcomes, we conducted sensitivity analyses that included modeling a dataset that only included mixed-breed dogs which was modeled in a mixed-effects framework that included breed as a random effect (SI Table 4, SI Figure 4).

### *SEM Confirmatory Factor analysis*

Confirmatory factor analysis was performed on a manually curated set of variables for use in the structural equation modeling analysis. This analysis was used to verify the structure of our observed set of variables. This was performed using the `cfa()` function under the Lavaan package (version 0.6-9) in R [48].

### *Model Specification*

The latent factor structure was defined based on known relationships in the literature studied primarily in humans ([49,50], SI Table 6). To specify the model for this study, we chose measured variables based on the following criteria, they: 1) had clear individual interpretability, 2) were likely to capture different aspects of the latent factor, 3) were continuously distributed, 4) did not have missing data, and 5) together maximized sample variance as recorded in factor analysis. Our latent factor of finance was informed using measures from the census median income, percent living below the poverty line, and percent of households making less than \$100,000 US per year as well as owner-reported annual income range. The latent factor of the local environment was informed by census-tract population density, neighborhood walk score, percent of individuals in the same household one year ago, and percent of the population that was born in the US. These variables were selected as measures that capture unique information about the dog's neighborhood environment. Lastly, our social environment latent variable was informed by the measurable variables of routine hours per day with dogs, children, and adults. We defined the structural equation model (SEM) using the Lavaan (version 0.6-9) package in R[48]; structural equation meta-models and model syntax were adapted from guidelines for a graph-theoretic implementation of structural equation modeling [51]. Prior to implementation of the SEM, thirteen variables were standardized to a mean of zero and standard deviation of 1 to account for scale differences among variables, and models were run using unweighted least squares. Two models were tested, with the difference being the addition of a pathway in model two from finance to environment. Fit for each of the two models was assessed through several metrics and associated cutoffs for "good" agreement between data and model: Root Mean Squared Error of Approximation (RMSEA, <0.08), Comparative Fit Index (CFI, >0.9), Standardized Root-Mean Squared Residual (SRMR, <0.08) and the Tucker-Lewis Index (TLI, >0.9) [52]. Comparative model fit was assessed using the corrected Akaike Information



Criterion (AICc). Confirmatory factor analysis on both latent and indicator variables was conducted using the `cfa` function in R. All structural model visualizations were generated using the `TidySEM` and `SEMplot` packages in R. We generated two models; the first model (M1) describes direct paths from the latent factor of finance, environment, and social interactions to our outcome of health (SI Figure 2). The second model (M2) describes all direct paths indicated in M1 but with the added indirect path between finance and environment (SI Figure 3). These two models were both hypothesized to reliably describe the system and were tested for model fit before interpreting the structural equation modeling analysis.

## Results

### **Owner survey data from a large cohort of companion dogs**

Our dataset included survey and health data from 21,410 dogs between the ages of 2 and 25.5 years old (Figure 1A), that weighed between 2-210 lbs (Figure 1B). We had a similar proportion of male (50.1%) and female dogs (49.8%; Fig 1C), as well as purebred (52.3%) and mixed-breed dogs (47.7%; Figure 1D). Given the low percentage of intact male and female dogs (4% and 2% respectively), we chose to include them in our analyses. Additional demographic information on the dogs and their owners can be found in Supplementary Information (Table S1, Figure S1).

### **Social, financial, and demographic factors**

We identified 5 factors, based on parallel analysis (see Methods), that collectively explained 33.7% of the variance in responses to the 33 survey questions, and then labeled the factors based on the survey questions that primarily contributed to each factor (loading  $>|0.4|$ ): neighborhood stability (factor 1: 10.6% variance explained), total household income (factor 2: 9.7% variance explained), social-children (factor 3: 5.1% variance explained), social-animals (factor 4: 4.4% variance explained), and owner age (factor 5: 3.9% variance explained; Figure 2).

## **Household environment is associated with dog health, disease, and mobility**

### *Owner-reported Dog Health*

Owner-reported dog health was reported as an integer on a scale from 1 (“very poor health”) to 6 (“excellent health”). Since this is an unrefined measure of health, we first assessed its internal validity to capture dog health. As expected based on previous literature, older and heavier dogs were rated by their owners as being in the poorest health ( $\beta_{\text{age}} = -1.01$ ,  $p \leq 2 \times 10^{-16}$ ;  $\beta_{\text{weight}} = -0.13$ ,  $p = 6.45 \times 10^{-23}$ ; Figure 3, [53,54]).

We found that all five factors were significantly associated with health outcomes (controlling for dog age and weight). Higher income was associated with better health ( $\beta_{\text{income}} = 0.037$ ,  $p = 5.99 \times 10^{-3}$ ), indicating that owners of higher SES rated their dogs as being healthier. Dogs with other dogs and/or other pets in their house were also rated as significantly healthier than dogs with fewer household companions ( $\beta_{\text{social-animals}} = 0.044$ ,  $p = 1.20 \times 10^{-3}$ ; Figure 4A). Dogs who lived in households with a higher stability score were reported to be healthier than those that did not ( $\beta_{\text{stability}} = 0.044$ ,  $p = 1.07 \times 10^{-45}$ ; Figure 4A). Dogs that lived in households with more children were reported to be less healthy than those with fewer children ( $\beta_{\text{children}} = -0.041$ ,  $p = 2.72 \times 10^{-3}$ ; Figure 4A). Finally, older owners reported their dogs were in better health compared to younger owners ( $\beta_{\text{owner\_age}} = 0.23$ ,  $p = 5.60 \times 10^{-41}$ ; Figure 4A).

### *Disease Diagnoses*

We found similar associations between environmental factors and health when using an arguably more objective measure of health: the total number of disease diagnoses per dog over their lifetime (the total count of all reported diagnosed diseases; see methods). All five factors were significantly associated with disease diagnoses, controlling for age and weight. Specifically, income was positively associated with the number of reported diseases, while stability, time with other animals, owner age, and time with children were negatively associated with the number of diseases reported. Dogs in more stable households or with older owners were reported to have fewer diseases ( $\beta_{\text{stability}} = -0.03$ ,  $p \leq 2 \times 10^{-16}$ ;  $\beta_{\text{owner\_age}}$

= -0.16,  $p \leq 2 \times 10^{-16}$ ). For the two measures of social integration and connectedness, we found that dogs with more nonhuman companions as well as those with more children in the house had significantly fewer reported diseases ( $\beta_{\text{social-animals}} = -0.03$ ,  $p = 9.09 \times 10^{-6}$ ,  $\beta_{\text{social-children}} = -0.09$ ,  $p \leq 2 \times 10^{-16}$ ) (Figure 4B). Paradoxically, we observed a positive association between income and reported disease instances ( $\beta_{\text{income}} = 0.04$ ,  $p = 2.27 \times 10^{-7}$ ), which stood in contrast to the negative association between higher income and better owner-reported health.

### *Mobility*

Given that mobility declines with age and can be used as a marker for exposures that might alter the rate of aging, we investigated associations between the environmental factors and composite mobility score that captured measures of physical activity duration, intensity, and frequency [55–57]. In this metric a higher score reflected higher levels of activity and mobility (see methods for details on calculation of the composite mobility score). As expected, older dogs were less mobile ( $\beta_{\text{age}} = -0.12$ ,  $p \leq 2 \times 10^{-16}$ ; SI Table 3), and, when controlling for age, we found that dogs with higher mobility were in better health compared to less mobile dogs of the same age ( $\beta = 0.35$ ,  $p \leq 2 \times 10^{-16}$ ; SI Table 3). This suggests that, similar to humans, a more active lifestyle is associated with better health in dogs [58,59]. We next found that the dog's social environment—specifically stability, time spent with children and owner age—were significantly associated with dog mobility. Mobility was lower in dogs that lived in households with more children and higher for dogs that lived in households with older owners ( $\beta_{\text{children}} = -0.04$ ,  $p \leq 2 \times 10^{-16}$ ,  $\beta_{\text{owner age}} = 0.03$ ,  $p = 1.15 \times 10^{-14}$ , respectively), while it was higher in dogs in households that were more stable ( $\beta_{\text{stability}} = 4.55 \times 10^{-3}$ ,  $p = 4.52 \times 10^{-12}$ ; Figure 4C, SI Table 3).

### **Effects of environmental determinants of health can vary across the lifespan**

To investigate whether associations between these factors and health, disease, and mobility vary across the lifespan, we extended each regression model by including an interaction term between each environmental factor and age (SI Figure 4). We found five significant interactions between dog age and our environmental factors on owner-reported dog health (SI Table 3). The positive association between household stability and dog health

was strongest in the younger dogs, and weakest in older dogs (Figure 5A,  $\beta = -0.03$ ,  $p = 5.33 \times 10^{-17}$ ). Time spent with other animals was more strongly associated with health when the dog was older rather than younger (Figure 5B,  $\beta = 0.03$ ,  $p = 4.18 \times 10^{-2}$ ). We also found that owner age was more impactful for younger dogs compared to older dogs (Figure 5C,  $\beta = -0.18$ ,  $p = 5.91 \times 10^{-26}$ ). Lastly, for older dogs, increased time spent with children was more impactful to overall general health than for younger dogs (SI Figure 6,  $\beta = -0.04$ ,  $p = 1.01 \times 10^{-2}$ ). We saw no significant interactions between environmental factors and age on disease instances.

We did find a significant interaction between dog age and four environmental factors on mobility. First, a significant interaction between neighborhood stability and dog age on mobility revealed that neighborhood stability is most strongly associated with the mobility of younger dogs compared to older dogs ( $\beta = -0.002$ ,  $p = 6.0 \times 10^{-4}$ ; SI Figure 5A; SI Table 3). Second, the effect of spending time with other animals on mobility significantly interacted with dog age, where more time was considerably more important for the mobility of older dogs ( $\beta = -0.01$ ,  $p = 8.2 \times 10^{-4}$ , SI Figure 5B; SI Table 3). Third, the effect of spending time with children on mobility significantly interacted with dog age, where more time with children was more important for the mobility of older dogs ( $\beta = -0.01$ ,  $p = 0.01$ , SI Figure 5C; SI Table 3). Lastly, having an older owner had a stronger effect on a younger dog's mobility ( $\beta = -0.01$ ,  $p = 7.9 \times 10^{-5}$ , Figure SI 5D).

### **Pathways linking social-environmental variables and owner-reported dog health.**

Environmental factors could have both direct and indirect effects on health. To examine these putative paths, we took an orthogonal modeling approach, structural equation modeling (SEM), which can capture more complex relationships among the environmental variables and health. This approach can have more flexibility, the capacity to represent indirect relationships, and supports integration of results into broader theory [51]. To do so, we first organized our variables into latent factors that represent concepts known to fundamentally influence mammalian health and that were chosen to maximize the interpretability of the latent factors [60]. We chose the variables (i.e., survey items) that were identified as most informative in our exploratory factor analysis described above. The latent factor of health was informed by the same variables that we used as outcome variables in our linear models: health, disease instances, and mobility.

Both showed good model fit according to established parameters [52] (M1: CFI = 0.84, RMSEA = 0.06, SRMR = 0.05, TLI = 0.80 ; M2: CFI = 0.84, RMSEA= 0.06, SRMR = 0.05, TLI = 0.80). Here we will present results for M2, which includes a pathway from the financial factor to the environmental factor, which is a better representation of the system given the established literature [61,62]. In humans we know that the quality of the environment is influenced by the finances of the household thus providing support for evaluating M2 as the more generalizable model [63].

As with our linear models, we found evidence for all three latent variables associated with the latent factor capturing health. However, the relative importance of the latent factors varied: the effect of the social factor was ~5x larger than the effect of finance and the local environment on our health outcome ( $\beta_{\text{Social}} = 0.12$  ,  $\beta_{\text{Finance}} = 0.02$ ,  $\beta_{\text{Environment}} = -0.02$ ). In line with our linear models, we found that increased income and more social interaction with humans and other dogs both predicted better health. In contrast, a more urban environment with less stability was associated with worse health, pointing to the complex relationship between overall health and the environment. These data capture a previously hypothesized indirect pathway from finance through the local environment that our linear models failed to identify, suggesting that the local environment might be an important mediator between finance and health and supporting the use of SEM to model these complex systems.

## Discussion

Using the largest observational dataset of companion dog environment and health, we identified measures of the social and lived environment that were linked to age-associated changes in health, mobility, and disease. Unsurprisingly, age was the strongest predictor of health deficits. However, not all dogs exhibited the same age-related changes in health, disease, and mobility, and some of this inter-individual variation was linked to the environment. Health, mobility, and disease measures were also associated with variation in household social environmental determinants. We also identified key changes in the environment that were associated with the health and mobility of dogs in an age-dependent manner within this cross-sectional dataset.

Similar to humans, we found that measures of income and social connectedness (i.e., presence of more potential social companions) predicted better health in dogs, although dogs from wealthier owners had more diagnosed diseases [64]. This counterintuitive association does not necessarily mean that dogs of wealthier owners had more diseases, especially given

the fact that dogs from higher SES households were reported as healthier by their owners. Rather, this finding points to the role that finance plays in the opportunities for disease diagnosis, while also generating new questions on the directionality of the relationship between income and health as well as what we ultimately consider “health” and how we diagnose diseases [60]. Dogs who live in households with wealthier owners might seek veterinary care more frequently and have the funds to pay for additional tests, resulting in more disease diagnoses even though the diseases are unlikely to be more prevalent in higher SES households. Owners may perceive their dogs to be in good health because they have more information and are more confident that any disease would have been detected and treated or managed. Dogs who lived in households with other pets had higher reported health scores and fewer disease diagnoses than dogs who had fewer household companions, suggesting that this social enrichment confers health benefits, which outweighed the effects of finances five-fold (Fig 6). These other companions are often other dogs—of homes with other animals present, 70% of them had another dog—but could be other pets (e.g., cats, SI Table 7). This suggests that similar to humans, increased social enrichment in companion dogs presents health benefits and also provides an example of an easily modifiable environmental intervention to improve dog welfare. Supporting the hypothesis from literature that shows interspecies companionship lowered frailty among older humans and may point to an effective intervention to improve the health of aging adults[65]. Dogs who lived in households with more children had fewer diseases, but tended to be in poorer health. This observation could be due to modified resource allocation (e.g., time and money) in households with more children. Mobility was also associated with environmental factors and showed unique age dependencies. This recapitulates what we see in humans, where individuals who live in lower income neighborhoods and experience adverse community-level factors, such as decreased access to green space, unequal resource distribution and access (e.g. healthcare or food), have poorer health outcomes [66,67].

We found that owner age was significantly associated with our health-related outcomes, but why this pattern occurs is unknown. One possible mechanism is the link between owner age and dog activity levels, where a recent analysis in the DAP found that older owners had more active dogs, which is linked to better health. Further answers to this question could elucidate the complex relationships between owner age and our other factors and their associations with health [68]. This suggests that as the DAP continues, investigators

will need to pay particular attention to owner age in our dataset, as it has a potential association with physical activity, cognitive function, and now perceived health.

An important outstanding question is identifying the path through which the social environment impacts health. In humans, it has been hypothesized (and there is some support for the fact) that exposure to psychosocial stressors from unpredictable environments (e.g., work-related stress, low SES, unpredictable environments, and low social integration) lastingly alter the stress-response pathway and, consequently, immune function and health [69]. Our findings fit within this framework and show that the negative health effects of unpredictable social environments can extend to companion dogs. However, more work is needed to identify whether or not the physiological pathways (e.g., stress pathway) are shared among these two cohabitating species.

We also unexpectedly found that younger dogs with older owners experienced better health outcomes compared to older dogs who had older owners. These data suggest the importance of understanding how age impacts lifestyle, which could have knock-on effects for companions and companion animals. Specifically, the human social environment might contribute to dog health indirectly through variables like owner age that affect how often a dog might engage in social enrichment activities due to their owner's activity budget and lifestyle. We also acknowledge that the distribution of dog age is highly non-random with respect to owner age, which may explain some of this association. We found that younger dogs who lived in less stable households were in worse health than older dogs who lived in stable homes, suggesting that, similar to humans, environmental adversity plays a particularly important role in early life [70–72]. Older dogs who lived in higher-income households were found to be in better health compared to older dogs in lower-income households, while the health of younger dogs was less impacted by income variation, suggesting that dogs may be more resilient to variation in this form of adversity earlier in life [73,74]. Younger dogs' mobility experienced a larger age-associated increase when living in more stable neighborhoods compared to older dogs, suggesting that the benefits of living in a more stable household, while important across the lifespan, is more impactful in earlier life. Older dogs who lived in less stable neighborhoods were much less mobile than older dogs in more stable neighborhoods, suggesting that, similar to older people in lower SES neighborhoods, dogs experience a reduction in activity that could ultimately be linked with poorer health outcomes [75,76].

Our environmental and health measures are not independent of one another, suggesting a more complex network of paths among variables and their association with health outcomes. We used SEM to capture direct and indirect effects of each variable in one model. The results of our SEM supported the independent models of each health variable, but allowed us to propose a holistic model for the environmental impacts on companion dog health and aging that is generalizable to similar work done on humans. The model points to direct effects of finance and the local environment on dog health, in addition to an indirect contribution of finance on health through its effect on the local environment. This indirect path is supported in the human literature, where we see that SES dictates where a person will live and thus their environment as well as access to healthcare and both of these contribute to both positive and negative health outcomes [68,77]. This indicates the latent structures affecting health outcomes in dogs and humans are similar, and may be a product of aspects of modern society as well as the deeper evolutionary history of dogs being selectively bred alongside humans for thousands of years.

While the strengths of this dataset include the large sample size (n= 21,410) of the study cohort and the breadth of the HLES survey data, there are some limitations. First, our dataset comprises both mixed-breed and purebred dogs, and genetic background and dog breed are both known to be associated with dog health and disease [20,53]. However, our results remained largely consistent when we examined pure and mixed breed dogs independently, as well as when we controlled for individual breeds based on owner-reports (SI Table 4). The Dog Aging Project is currently sequencing the genomes of 10,000 dogs in the project [16], which will allow us to more accurately control for genetic background as well as identify putative gene-by-environment interactions that impact health, disease, or mobility.

Second, all data in the HLES are owner-reported and can thus be affected by subjective error, bias, and/or misinterpretation of survey questions. Further investigation of biases in owner reported responses will be needed to verify our findings in the future. The strong association between dog age and all three of our outcome measures—health, disease incidence, and mobility—suggests that the survey is an instrument that, on average, accurately captures the variables we cannot objectively measure. All measures could be improved by an owner-independent measure or veterinarian-informed measure of frailty and health. In future research, we will draw on electronic veterinary medical records, molecular and



immunological measures, and at-home physical tests to generate more accurate measures of health and frailty in the companion dog.

Third, there is a limitation due to the average household income being well above the U.S median income (US median: \$67,521 in 2020, DAP median: \$100,000-\$119,999, [78]) and biases our knowledge away from lower-earning households. This underscores the need to recruit a diverse set of human participants to better account for the variation of environmental exposures a dog is subject to [16].

Fourth, given the observational nature of the data, we were not able to determine causality or directionality among variables; we instead drew a causal structure in the SEM based on a critical evaluation of the literature (SI Table 6). For instance, owners who have dogs in poorer health might limit exposures to demanding activities or social enrichment compared to owners who have dogs who are in excellent health, which would also lead to an association between health and social companionship and activity.

Given the above limitations, the results of this analysis should not be used to influence medical decisions, behavioral interventions, or policy changes related to the care of companion dogs, but rather encourage the continued investigation by veterinary professionals and research scientists on best practices for increasing companion welfare. Some of these relationships can be evaluated in the future with randomized, controlled experiments.

Overall, our study provides further evidence for the strong link between the social environment and health outcomes that reflects what is known for humans. From this knowledge we propose the companion dog to be a novel model for human aging. This highlights the need for more attention to the role of the social environment on health and disease and continued investigation of how each environmental factor can contribute to more years of healthy living (i.e., “healthspan”) in both companion dogs and humans. A holistic understanding of environmental effects on both humans and dogs may offer insight to what factors are important for health-independent of confounding factors that complicate human studies (e.g., smoking). Results highlight the need to study how differences in resource allocation among households contributes to health and wellbeing of all members of the household. A future use of longitudinal data will allow stronger inferential capacity to identify putative causal factors that affect trajectories of health and aging in both dogs and humans, driving new insights into what factors promote healthy aging and potential lifestyle modifications that can increase time spent free of disease.

## Author Contributions

All authors contributed to writing – review & editing. B.M.M.: conceptualization, methodology, data curation, writing – original draft, formal analysis, visualization, and project administration. L.B.: conceptualization, methodology, data curation, formal analysis, writing – original draft and visualization. K.J.: data curation, formal analysis, and visualization. G.A.D.: data curation, formal analysis, methodology and visualization, manuscript editing. A.R.: manuscript curation, drafting and editing. S.S.: conceptualization. D.C.: data curation. M.D.: data curation. DAP consortium: resources and manuscript editing. N.S.M.: conceptualization, methodology, writing – original draft, funding acquisition, and project administration.

### Dog Aging Project Consortium Authors (as of March 2022)

Joshua M. Akey<sup>1</sup>, Brooke Benton<sup>2</sup>, Elhanan Borenstein<sup>3</sup>, Marta G. Castelhana<sup>4</sup>, Amanda E. Coleman<sup>5</sup>, Kate E. Creevy<sup>6</sup>, Kyle Crowder<sup>7</sup>, Matthew D. Dunbar<sup>8</sup>, Virginia R. Fajt<sup>9</sup>, Annette L. Fitzpatrick<sup>10</sup>, Unity Jeffery<sup>11</sup>, Erica C Jonlin<sup>12</sup>, Matt Kaeberlein<sup>13</sup>, Elinor K. Karlsson<sup>14</sup>, Kathleen F. Kerr<sup>15</sup>, Jonathan M. Levine<sup>16</sup>, Jing Ma<sup>17</sup>, Robyn L McClelland<sup>18</sup>, Daniel E.L. Promislow<sup>19</sup>, Audrey Ruple<sup>20</sup>, Stephen M. Schwartz<sup>21</sup>, Sandi Shrager<sup>22</sup>, Noah Snyder-Mackler<sup>23</sup>, Katherine Tolbert<sup>24</sup>, Silvan R. Urfer<sup>25</sup>, Benjamin S. Wilfond<sup>26</sup>

<sup>1</sup> Lewis-Sigler Institute for Integrative Genomics, Princeton University, Princeton, NJ, USA

<sup>2</sup> Department of Laboratory Medicine and Pathology, University of Washington School of Medicine, Seattle, WA, USA

<sup>3</sup> Department of Clinical Microbiology and Immunology, Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel

<sup>4</sup> Cornell Veterinary Biobank, College of Veterinary Medicine, Cornell University, Ithaca, NY, USA

<sup>5</sup> Department of Small Animal Medicine and Surgery, College of Veterinary Medicine, University of Georgia, Athens, GA, USA

<sup>6</sup> Department of Small Animal Clinical Sciences, Texas A&M University College of Veterinary Medicine & Biomedical Sciences, College Station, TX, USA

<sup>7</sup> Department of Sociology, University of Washington, Seattle, WA, USA

<sup>8</sup> Center for Studies in Demography and Ecology, University of Washington, Seattle, WA, USA

<sup>9</sup> Department of Veterinary Physiology and Pharmacology, Texas A&M University College of Veterinary Medicine & Biomedical Sciences, College Station, TX, USA

<sup>10</sup> Department of Family Medicine, University of Washington, Seattle, WA, USA

- <sup>11</sup> Department of Veterinary Pathobiology, Texas A&M University College of Veterinary Medicine & Biomedical Sciences, College Station, TX, USA
- <sup>12</sup> Department of Laboratory Medicine and Pathology, University of Washington School of Medicine, Seattle, WA, USA
- <sup>13</sup> Department of Laboratory Medicine and Pathology, University of Washington School of Medicine, Seattle, WA, USA
- <sup>14</sup> Bioinformatics and Integrative Biology, University of Massachusetts Chan Medical School, Worcester, MA, USA
- <sup>15</sup> Department of Biostatistics, University of Washington, Seattle, WA, USA
- <sup>16</sup> Department of Small Animal Clinical Sciences, Texas A&M University College of Veterinary Medicine & Biomedical Sciences, College Station, TX, USA
- <sup>17</sup> Division of Public Health Sciences, Fred Hutchinson Cancer Research Center, Seattle, WA, USA
- <sup>18</sup> Department of Biostatistics, University of Washington, Seattle, WA, USA
- <sup>19</sup> Department of Laboratory Medicine and Pathology, University of Washington School of Medicine, Seattle, WA, USA
- <sup>20</sup> Department of Population Health Sciences, Virginia-Maryland College of Veterinary Medicine, Virginia Tech, Blacksburg, VA, USA
- <sup>21</sup> Epidemiology Program, Fred Hutchinson Cancer Research Center, Seattle, WA, USA
- <sup>22</sup> Collaborative Health Studies Coordinating Center, Department of Biostatistics, University of Washington, Seattle, WA, USA
- <sup>23</sup> School of Life Sciences, Arizona State University, Tempe, AZ, USA
- <sup>24</sup> Department of Small Animal Clinical Sciences, Texas A&M University College of Veterinary Medicine & Biomedical Sciences, College Station, TX, USA
- <sup>25</sup> Department of Laboratory Medicine and Pathology, University of Washington School of Medicine, Seattle, WA, USA
- <sup>26</sup> Treuman Katz Center for Pediatric Bioethics, Seattle Children's Research Institute, Seattle, WA, USA

### **Acknowledgments**

The authors thank Dog Aging Project participants, their dogs, and community veterinarians for their important contributions. The Dog Aging Project is supported by U19 grant AG057377 from the National Institute on Aging, a part of the National Institutes of Health, and by private donations. BMM is supported by a Graduate Research Fellowship from the National Science Foundation. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

## References

1. Snyder-Mackler N, Burger JR, Gaydosh L, Belsky DW, Noppert GA, Campos FA, et al. Social determinants of health and survival in humans and other animals. *Science*. 2020;368. doi:10.1126/science.aax9553
2. Weyers S, Dragano N, Möbus S, Beck E-M, Stang A, Möhlenkamp S, et al. Low socioeconomic position is associated with poor social networks and social support: results from the Heinz Nixdorf Recall Study. *Int J Equity Health*. 2008;7: 13.
3. Holt-Lunstad J, Smith TB, Layton JB. Social relationships and mortality risk: a meta-analytic review. *PLoS Med*. 2010;7: e1000316.
4. Algren MH, Ekholm O, Nielsen L, Ersbøll AK, Bak CK, Andersen PT. Social isolation, loneliness, socioeconomic status, and health-risk behaviour in deprived neighbourhoods in Denmark: A cross-sectional study. *SSM - Population Health*. 2020. p. 100546. doi:10.1016/j.ssmph.2020.100546
5. Hawkey LC, Cole SW, Capitanio JP, Norman GJ, Cacioppo JT. Effects of social isolation on glucocorticoid regulation in social mammals. *Horm Behav*. 2012;62: 314–323.
6. Sullens DG, Gilley K, Jensen K, Vichaya E, Dolan SL, Sekeres MJ. Social isolation induces hyperactivity and exploration in aged female mice. *PLoS One*. 2021;16: e0245355.
7. Razzoli M, Nyuyki-Dufe K, Gurney A, Erickson C, McCallum J, Spielman N, et al. Social stress shortens lifespan in mice. *Aging Cell*. 2018;17: e12778.
8. Scott SB, Sliwinski MJ, Blanchard-Fields F. Age differences in emotional responses to daily stress: the role of timing, severity, and global perceived stress. *Psychol Aging*. 2013;28: 1076–1087.
9. Sarkar C, Webster C. Urban environments and human health: current trends and future directions. *Current Opinion in Environmental Sustainability*. 2017;25: 33–44.
10. Vlahov D, Freudenberg N, Proietti F, Ompad D, Quinn A, Nandi V, et al. Urban as a Determinant of Health. *Journal of Urban Health*. 2007. pp. 16–26. doi:10.1007/s11524-007-9169-3
11. Zaninotto ASA. Lower socioeconomic status and the acceleration of aging: An outcome-wide analysis. *PNAS*. 2020. doi:10.1073/pnas.1915741117
12. Zeng Y, Gu D, Purser J, Hoenig H, Christakis N. Associations of environmental factors with elderly health and mortality in China. *Am J Public Health*. 2010;100: 298–305.
13. Hadley MB, Nalini M, Adhikari S, Szymonifka J, Etemadi A, Kamangar F, et al. Spatial environmental factors predict cardiovascular and all-cause mortality: Results of the SPACE study. *PLoS One*. 2022;17: e0269650.
14. Ruple A, MacLean E, Snyder-Mackler N, Creevy KE, Promislow D. Dog Models of Aging. *Annu Rev Anim Biosci*. 2022;10: 419–439.

15. Hoffman JM, Creevy KE, Franks A, O'Neill DG, Promislow DEL. The companion dog as a model for human aging and mortality. *Aging Cell*. 2018;17: e12737.
16. Creevy KE, Akey JM, Kaeberlein M, Promislow DEL, Dog Aging Project Consortium. An open science study of ageing in companion dogs. *Nature*. 2022;602: 51–57.
17. Why is the dog an ideal model for aging research? *Exp Gerontol*. 2015;71: 14–20.
18. The companion dog as a unique translational model for aging. *Semin Cell Dev Biol*. 2017;70: 141–153.
19. Dias-Pereira P. Morbidity and mortality in elderly dogs – a model for human aging. *BMC Vet Res*. 2022;18: 1–8.
20. Fleming JM, Creevy KE, Promislow DEL. Mortality in north american dogs from 1984 to 2004: an investigation into age-, size-, and breed-related causes of death. *J Vet Intern Med*. 2011;25: 187–198.
21. Vaysse A, Ratnakumar A, Derrien T, Axelsson E, Rosengren Pielberg G, Sigurdsson S, et al. Identification of genomic regions associated with phenotypic variation between dog breeds using selection mapping. *PLoS Genet*. 2011;7: e1002316.
22. Vonholdt BM, Pollinger JP, Lohmueller KE, Han E, Parker HG, Quignon P, et al. Genome-wide SNP and haplotype analyses reveal a rich history underlying dog domestication. *Nature*. 2010;464: 898–902.
23. Shearin AL, Ostrander EA. Leading the way: canine models of genomics and disease. *Dis Model Mech*. 2010;3: 27–34.
24. Karlsson EK, Lindblad-Toh K. Leader of the pack: gene mapping in dogs and other model organisms. *Nat Rev Genet*. 2008;9: 713–725.
25. Whitson HE, Cohen HJ, Schmader KE, Morey MC, Kuchel G, Colon-Emeric CS. Physical Resilience: Not Simply the Opposite of Frailty. *Journal of the American Geriatrics Society*. 2018. pp. 1459–1461. doi:10.1111/jgs.15233
26. Whitson HE, Duan-Porter W, Schmader KE, Morey MC, Cohen HJ, Colón-Emeric CS. Physical Resilience in Older Adults: Systematic Review and Development of an Emerging Construct. *J Gerontol A Biol Sci Med Sci*. 2016;71: 489–495.
27. Candela Andrade M, Slunsky P, Klass LG, Brunnberg L. Risk factors and long-term surgical outcome of patellar luxation and concomitant cranial cruciate ligament rupture in small breed dogs. *Vet Med*. 2020;65: 159–167.
28. Allenspach K, Wieland B, Gröne A, Gaschen F. Chronic enteropathies in dogs: Evaluation of risk factors for negative outcome. *J Vet Intern Med*. 2007;21: 700–708.
29. Gillman MW. Developmental origins of health and disease. *The New England journal of medicine*. 2005. pp. 1848–1850.
30. Vargas J, Junco M, Gomez C, Lajud N. Early Life Stress Increases Metabolic Risk, HPA Axis Reactivity, and Depressive-Like Behavior When Combined with

Postweaning Social Isolation in Rats. *PLoS One*. 2016;11: e0162665.

31. Pesonen A-K, Räikkönen K, Heinonen K, Kajantie E, Forsén T, Eriksson JG. Depressive symptoms in adults separated from their parents as children: a natural experiment during World War II. *Am J Epidemiol*. 2007;166: 1126–1133.
32. Heim C, Newport DJ, Mletzko T, Miller AH, Nemeroff CB. The link between childhood trauma and depression: insights from HPA axis studies in humans. *Psychoneuroendocrinology*. 2008;33: 693–710.
33. Pienaar AE. The association between under-nutrition, school performance and perceptual motor functioning in first-grade South African learners: The North-West Child Health Integrated with Learning and Development study. *Health Saf Code Annot State Calif Adopt April 7 1939 Calif*. 2019;24: 1046.
34. Houpt KA, Zicker S. Dietary effects on canine and feline behavior. *Vet Clin North Am Small Anim Pract*. 2003;33: 405–16, vii–viii.
35. Bisset ES, Howlett SE. The biology of frailty in humans and animals: Understanding frailty and promoting translation. *Aging Med (Milton)*. 2019;2: 27–34.
36. Chen FL, Ullal TV, Graves JL, Ratcliff ER, Naka A, McKenzie B, et al. Evaluating instruments for assessing healthspan: a multi-center cross-sectional study on health-related quality of life (HRQL) and frailty in the companion dog. *bioRxiv*. 2022. p. 2022.07.21.500746. doi:10.1101/2022.07.21.500746
37. Rantakokko M, Mänty M, Rantanen T. Mobility decline in old age. *Exerc Sport Sci Rev*. 2013;41: 19–25.
38. Morgan EM, Heseltine JC, Levine GJ, Promislow DEL, Creevy KE. Evaluation of a low-technology system to obtain morphological and mobility trial measurements in dogs and investigation of potential predictors of canine mobility. *Am J Vet Res*. 2019;80: 670–679.
39. Kaeberlein M, Creevy KE, Promislow DEL. The dog aging project: translational geroscience in companion animals. *Mammalian Genome*. 2016. pp. 279–288. doi:10.1007/s00335-016-9638-7
40. Creevy KE, Akey JM, Kaeberlein M, Promislow DEL. Author Correction: An open science study of ageing in companion dogs. *Nature*. 2022;608. doi:10.1038/s41586-022-05179-x
41. Harris PA, Taylor R, Minor BL, Elliott V, Fernandez M, O’Neal L, et al. The REDCap consortium: Building an international community of software platform partners. *J Biomed Inform*. 2019;95: 103208.
42. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)--a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform*. 2009;42: 377–381.
43. Creevy KE, Grady J, Little SE, Moore GE, Strickler BG, Thompson S, et al. 2019

- AAHA Canine Life Stage Guidelines. *J Am Anim Hosp Assoc.* 2019;55: 267–290.
44. Bray EE, Zheng Z, Tolbert MK, McCoy BM, Dog Aging Project Consortium, Kaeberlein M, et al. Once-daily feeding is associated with better health in companion dogs: results from the Dog Aging Project. *Geroscience.* 2022;44: 1779–1790.
  45. Fabrigar LR, Wegener DT, MacCallum RC, Strahan EJ. Evaluating the use of exploratory factor analysis in psychological research. *Psychological Methods.* 1999. pp. 272–299. doi:10.1037/1082-989x.4.3.272
  46. Website. Available: Revelle W (2021). psych: Procedures for Psychological, Psychometric, and Personality Research. Northwestern University, Evanston, Illinois. R package version 2.1.9, <https://CRAN.R-project.org/package=psych>.
  47. Kraus C, Pavard S, Promislow DEL. The size-life span trade-off decomposed: why large dogs die young. *Am Nat.* 2013;181: 492–505.
  48. Rosseel Y. lavaan: AnRPackage for Structural Equation Modeling. *Journal of Statistical Software.* 2012. doi:10.18637/jss.v048.i02
  49. Marmot M, Wilkinson RG. Social determinants of health in older age. *Social Determinants of Health.* 2005. pp. 267–296. doi:10.1093/acprof:oso/9780198565895.003.13
  50. Jian Wang LG. Effects of Socioeconomic Status on Physical and Psychological Health: Lifestyle as a Mediator. *Int J Environ Res Public Health.* 2019;16. doi:10.3390/ijerph16020281
  51. Grace JB, Schoolmaster DR, Guntenspergen GR, Little AM, Mitchell BR, Miller KM, et al. Guidelines for a graph-theoretic implementation of structural equation modeling. *Ecosphere.* 2012. p. art73. doi:10.1890/es12-00048.1
  52. Hu L, Bentler PM. Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Struct Equ Modeling.* 2009 [cited 13 Dec 2021]. doi:10.1080/10705519909540118
  53. Yordy J, Kraus C, Hayward JJ, White ME, Shannon LM, Creevy KE, et al. Body size, inbreeding, and lifespan in domestic dogs. *Conserv Genet.* 2020;21. doi:10.1007/s10592-019-01240-x
  54. German AJ. The Growing Problem of Obesity in Dogs and Cats. *J Nutr.* 2006;136: 1940S–1946S.
  55. Ferrucci L, Cooper R, Shardell M, Simonsick EM, Schrack JA, Kuh D. Age-Related Change in Mobility: Perspectives From Life Course Epidemiology and Geroscience. *J Gerontol A Biol Sci Med Sci.* 2016;71: 1184.
  56. Guralnik JM, Ferrucci L, Simonsick EM, Salive ME, Wallace RB. Lower-extremity function in persons over the age of 70 years as a predictor of subsequent disability. *N Engl J Med.* 1995;332. doi:10.1056/NEJM199503023320902
  57. Studenski S, Perera S, Patel K, Rosano C, Faulkner K, Inzitari M, et al. Gait Speed and

Survival in Older Adults. *JAMA*. 2011;305: 50.

58. Powell KE, Paluch AE, Blair SN. Physical Activity for Health: What Kind? How Much? How Intense? On Top of What? *Annual Review of Public Health*. 2011. pp. 349–365. doi:10.1146/annurev-publhealth-031210-101151
59. Lee IM, Shiroma EJ, Lobelo F, Puska P, Blair SN, Katzmarzyk PT. Effect of physical inactivity on major non-communicable diseases worldwide: an analysis of burden of disease and life expectancy. *Lancet*. 2012;380. doi:10.1016/S0140-6736(12)61031-9
60. Daly M, Boyce C, Wood A. A social rank explanation of how money influences health. *Health Psychol*. 2015;34: 222–230.
61. Barrington WE. *Neighborhood Environment, Stress, and Obesogenic Behaviors Among Adults*. 2012.
62. The stress process in neighborhood context. *Health Place*. 2000;6: 287–299.
63. Evans and GW, Kantrowitz E. *Socioeconomic Status and Health: The Potential Role of Environmental Risk Exposure*. 2003 [cited 19 Jan 2022]. doi:10.1146/annurev.publhealth.23.112001.112349
64. Simandan D. Rethinking the health consequences of social class and social mobility. *Soc Sci Med*. 2018;200: 258–261.
65. Taniguchi Y, Seino S, Nishi M, Tomine Y, Tanaka I, Yokoyama Y, et al. Association of Dog and Cat Ownership with Incident Frailty among Community-Dwelling Elderly Japanese. *Sci Rep*. 2019;9: 1–7.
66. Diez Roux AV, Mair CF. Neighborhoods and health. *Ann N Y Acad Sci*. 2010;1186: 125–145.
67. Grafova IB, Freedman VA, Kumar R, Rogowski J. Neighborhoods and Obesity in Later Life. *American Journal of Public Health*. 2008. pp. 2065–2071. doi:10.2105/ajph.2007.127712
68. Balia S, Jones AM. Mortality, lifestyle and socio-economic status. *Journal of Health Economics*. 2008. pp. 1–26. doi:10.1016/j.jhealeco.2007.03.001
69. Rohleder N. Stimulation of Systemic Low-Grade Inflammation by Psychosocial Stress. *Psychosom Med*. 2014;76: 181.
70. Nelson CA, Scott RD, Bhutta ZA, Harris NB, Danese A, Samara M. Adversity in childhood is linked to mental and physical health throughout life. *BMJ*. 2020;371: m3048.
71. Suglia SF, Campo RA, Brown AGM, Stoney C, Boyce CA, Appleton AA, et al. Social Determinants of Cardiovascular Health: Early Life Adversity as a Contributor to Disparities in Cardiovascular Diseases. *J Pediatr*. 2020;219: 267–273.
72. McCrory C, Dooley C, Layte R, Kenny RA. The lasting legacy of childhood adversity for disease risk in later life. *Health Psychol*. 2015;34: 687–696.



73. Belsky J. The Differential Susceptibility Hypothesis: Sensitivity to the Environment for Better and for Worse. *JAMA pediatrics*. 2016. pp. 321–322.
74. Esther Nederhof MVS. Mismatch or cumulative stress: Toward an integrated hypothesis of programming effects. *Physiol Behav*. 2012;106: 691–700.
75. Talaei M, Rabiei K, Talaei Z, Amiri N, Zolfaghari B, Kabiri P, et al. Physical activity, sex, and socioeconomic status: A population based study. *ARYA Atheroscler*. 2013;9: 51.
76. Powell LM, Slater S, Chaloupka FJ, Harper D. Availability of Physical Activity–Related Facilities and Neighborhood Demographic and Socioeconomic Characteristics: A National Study. *Am J Public Health*. 2006;96: 1676.
77. Contoyannis P, Jones AM. Socio-economic status, health and lifestyle. *Journal of Health Economics*. 2004. pp. 965–995. doi:10.1016/j.jhealeco.2004.02.001
78. US Census Bureau. Income and Poverty in the United States: 2020. 2021 [cited 16 Dec 2021]. Available: <https://www.census.gov/library/publications/2021/demo/p60-273.html>

Accepted Manuscript

**Figure 1. Demographic characteristics of dogs included in this study.** Distribution of (a) age and (b) weight of dogs included in this study. The sample was roughly balanced by (c) sex and (d) mixed vs. purebred ancestry. Data are from the Dog Aging Project Health and Life Experience (HLES) Survey, 2019-2020.

**Figure 2. Five factors capture much of the social environmental variation in the Dog Aging Project cohort.** (a) Exploratory factor analysis revealed five factors that together explained 33.7% of the variance in survey responses. We named each factor based on the survey questions that loaded the strongest onto each factor (loading > |0.4|). (b) Distribution of factor scores for individual dogs for each of the 5 factors in our sample.


**Figure 3. Owner-reported health is worse in older dogs.** (a) Dog age was significantly associated with owner-reported health, such that older dogs were reported to be in poorer health compared to younger dogs controlling for dog weight ( $\beta_{\text{age}} = -0.421$ ,  $p \leq 2 \times 10^{-16}$ ).

**Figure 4. Social environment factors are associated with health, disease, and mobility in companion dogs.** Effect sizes of each environmental factor on (a) owner-reported dog health (ordinal model), (b) disease prevalence (cumulative number of diseases reported) (generalized linear model with Poisson link function), and (c) mobility (composite measure of activity level) (linear model). Points depict the effect sizes ( $\beta$ ) of each factor and lines represent the 95% confidence intervals.

**Figure 5. Age-dependent associations between environmental factors and health.** (a) Household stability was more strongly associated with health for younger dogs compared to older dogs. (b) Dog sociability was more strongly associated with health for older dogs compared to younger dogs. (c) The factor of owner age was more strongly associated with health for younger dogs compared to older dogs. (d) More time spent with children was more strongly associated with health for older dogs compared to younger dogs. All graphs show predicted trends using a linear model framework with 95% confidence intervals for a hypothetical population of dogs aged 4 (- 1 SD, lightest purple), 7.9 (mean age, intermediate purple), and 11.8 (+ 1 SD, darkest purple) years old.


**Figure 6. Structural equation modeling captures a complex network of environmental effects on health.** Structural equation model of how the social environment can influence general health. Path coefficients explain effect sizes and directionality of each major path. Latent factors (circles) are informed by measured variables (squares) that were chosen via factor analysis, and the direction of the effect of each measured variable on the latent variables is indicated next to or below the variable (+: positive association; -: negative association).

Figure 1




Accepted Manuscript

Figure 2




Accepted Manuscript

Figure 3




Accepted Manuscript

Figure 4




Accepted Manuscript

Figure 5



Accepted Manuscript

Figure 6



Accepted Manuscript